

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N	A2	(11) International Publication Number: WO 97/07198 (43) International Publication Date: 27 February 1997 (27.02.97)
(21) International Application Number: PCT/US96/12897 (22) International Filing Date: 8 August 1996 (08.08.96) (30) Priority Data: Not furnished 11 August 1995 (11.08.95) US (71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US). (72) Inventors: JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). MCCOY, John, M.; 63 Pine Ridge Road, Reading, MA 01867 (US). KELLEHER, Kerry; 50 Hurley Circle, Marlborough, MA 01752 (US). CARLIN, McKeough; 16 Chauncy Street #22, Cambridge, MA 02138 (US). (74) Agent: BROWN, Scott, A.; Genetics Institute, Inc., Legal Affairs, 87 CambridgePark Drive, Cambridge, MA 02140 (US).		(81) Designated States: AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: DNA SEQUENCES AND SECRETED PROTEINS ENCODED THEREBY (57) Abstract Novel polynucleotides and the proteins encoded thereby are disclosed.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DNA SEQUENCES AND SECRETED PROTEINS ENCODED THEREBY

5 This application claims priority from application Ser. No. 08/514,014, filed on August 11, 1995, which was converted to provisional application Ser. No. 60/_____ on July 19, 1996.

FIELD OF THE INVENTION

10 The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins.

BACKGROUND OF THE INVENTION

15 Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered factor (i.e., partial DNA/amino acid
20 sequence of the factor in the case of hybridization cloning; activity of the factor in the case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by
25 making available large numbers of DNA/amino acid sequences for factors that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these factors and the polynucleotides encoding them that the present invention is directed.

30

5

SUMMARY

In one embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 38 to nucleotide 1447;
- 10 (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:1 encoding a protein having biological activity;
- (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
- (d) a polynucleotide encoding a protein comprising a fragment of
15 the amino acid sequence of SEQ ID NO:2 having biological activity;
- (e) a polynucleotide which is an allelic variant of SEQ ID NO:1;
and
- (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

20

In another embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 52 to nucleotide 2034;
- (b) a polynucleotide comprising a fragment of the nucleotide
25 sequence of SEQ ID NO:3 encoding a protein having biological activity;
- (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity;
- 30 (e) a polynucleotide which is an allelic variant of SEQ ID NO:4;
and
- (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

35 In another embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 76 to nucleotide 474;

(b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:5 encoding a protein having biological activity;

10 (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;

(d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity;

(e) a polynucleotide which is an allelic variant of SEQ ID NO:5; and

15 (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

In another embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

20 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 67 to nucleotide 348;

(b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:7 encoding a protein having biological activity;

(c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:8;

25 (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity;

(e) a polynucleotide which is an allelic variant of SEQ ID NO:7; and

30 (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

In another embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 75 to nucleotide 356;

35 (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:9 encoding a protein having biological activity;

5 (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;

(d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity;

10 (e) a polynucleotide which is an allelic variant of SEQ ID NO:9; and

(f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

In another embodiment, the present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 86 to nucleotide 544;

(b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:11 encoding a protein having biological activity;

20 (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;

(d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity;

(e) a polynucleotide which is an allelic variant of SEQ ID NO:11; and

25 (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

In certain preferred embodiments, the polynucleotide is operably linked to an expression control sequence. The invention also provides a host cell, including bacterial, yeast, insect and mammalian cells, transformed with such polynucleotide compositions.

30 Processes are also provided for producing a protein, which comprise:

(a) growing a culture of the host cell transformed with such polynucleotide compositions in a suitable culture medium; and

(b) purifying the protein from the culture.

35 The protein produced according to such methods is also provided by the present invention.

5 Compositions comprising a protein biological activity are also disclosed. In preferred embodiments the protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- (b) fragments of the amino acid sequence of SEQ ID NO:2;
- 10 (c) the amino acid sequence of SEQ ID NO:4;
- (d) fragments of the amino acid sequence of SEQ ID NO:4;
- (e) the amino acid sequence of SEQ ID NO:6;
- (f) fragments of the amino acid sequence of SEQ ID NO:6;
- (g) the amino acid sequence of SEQ ID NO:8;
- 15 (h) fragments of the amino acid sequence of SEQ ID NO:8;
- (i) the amino acid sequence of SEQ ID NO:12; and
- (j) fragments of the amino acid sequence of SEQ ID NO:12;

the protein being substantially free from other mammalian proteins.

Such compositions may further comprise a pharmaceutically acceptable carrier.

20 Compositions comprising an antibody which specifically reacts with such protein are also provided by the present invention.

Methods are also provided for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition comprising a protein of the present invention and
25 a pharmaceutically acceptable carrier.

BRIEF DESCRIPTION OF FIGURES

Fig. 1 is an autoradiograph evidencing the expression of clone J5 in COS cells (indicated by arrows). J5 is processed into multiple bands, with the major band at
30 approximately 58 kD.

Fig. 2 is an autoradiograph evidencing the expression of clone L105 in COS cells (indicated by arrows).

Fig. 3 is an autoradiograph evidencing the expression of clone H174 in COS cells (indicated by arrows).

35 Fig. 4 is an autoradiograph evidencing the expression of clone B18 in COS cells (indicated by arrows).

5

DETAILED DESCRIPTIONISOLATED PROTEINS AND POLYNUCLEOTIDES

The sequence of a polynucleotide encoding one protein of the present invention is set forth in SEQ ID NO:1, with the coding region extending from nucleotides 38 to 1447. This polynucleotide has been identified as "clone J5". The amino acid sequence of the protein encoded by clone J5 is set forth in SEQ ID NO:2. Clone J5 was deposited with the American Type Culture Collection on August 11, 1995 and given the accession number ATCC 69885. SEQ ID NO:1 represents a spliced combination of sequence obtained from an isolated clone identified as "J5_3_fl", with additional 5' sequence obtained from a second double stranded clone. Clone J5 was isolated from a human activated peripheral blood mononuclear cell (PBMC) library using a trap which selects for nucleotides encoding secreted proteins; therefore, clone J5 does encode a secreted factor. J5 encodes a novel protein; BLASTN/BLASTX or FASTA searches revealed no exact sequence matches. However, a BLASTX search revealed homology between the J5 protein (in the approximate region of amino acids 62-129 of SEQ ID NO:2), epididymal apical proteins (including without limitation, epididymal apical protein I-precursor (*Macaca fascicularis*) (accession X66139)) and several snake venom haemorrhagic peptides (disintegrins) (including without limitation those assigned accession U01235-1237, X68251, and M89784). Analysis of the full-length J5 sequences revealed that the disintegrin domain was incomplete and that this clone did not contain an EGF-domain, as with some of the other disintegrin family members. J5 does contain a conserved metallo-proteinase domain. Based upon these homologies, J5 and these homologous proteins are expected to share at least some activities.

The sequence of a polynucleotide encoding another protein of the present invention is set forth in SEQ ID NO:3, with the coding region extending from nucleotides 52 to 2034. This polynucleotide has been identified as "clone J422". The amino acid sequence of the protein encoded by clone J422 is set forth in SEQ ID NO:4. Clone J422 was deposited with the American Type Culture Collection on August 11, 1995 and given the accession number ATCC 69884. SEQ ID NO:3 represents a spliced combination of sequence obtained from an isolated clone

5 identified as "J422_fl", with additional 5' sequence obtained from a second double stranded clone. Clone J422 was isolated from a human activated peripheral blood mononuclear cell (PBMC) library using a trap which selects for nucleotides encoding secreted proteins; therefore, clone J422 does encode a secreted factor. J422 encodes a novel protein; BLASTN/BLASTX or FASTA searches revealed no exact sequence
10 matches. However, a FASTA search revealed homology between the J422 protein (in the approximate region of amino acids 34-156 of SEQ ID NO:4) and a number of *Drosophila* leucine-rich repeat (LRR) proteins. Analysis of the full-length J422 sequences revealed that the conserved EGF-domain found in a number of LRR family members was not present in J422. Based upon these homologies, J422 and these
15 homologous proteins are expected to share at least some activities.

The sequence of a polynucleotide encoding another protein of the present invention is set forth in SEQ ID NO:5, with the coding region extending from nucleotides 76 to 474. This polynucleotide has been identified as "clone L105" The amino acid sequence of the protein encoded by clone L105 is set forth in SEQ ID
20 NO:6. Clone L105 was deposited with the American Type Culture Collection on August 11, 1995 and given the accession number ATCC 69883. Clone L105 was isolated from a murine adult thymus library using a trap which selects for nucleotides encoding secreted proteins; therefore, clone L105 does encode a secreted factor. L105 encodes a novel protein; BLASTN/BLASTX or FASTA searches revealed no exact
25 sequence matches. However, a BLASTX search revealed homology between the L105 protein (particularly in the approximate region of amino acids 73-91 of SEQ ID NO:6), various monocyte and other chemoattractant proteins (including without limitation those assigned accession M577441, X71087, X72308, X14768 and M24545) and a chicken (*Gallus gallus*) cytokine (accession L34553). Based upon
30 these homologies, L105 and these homologous proteins are expected to share at least some activities.

The sequence of polynucleotides encoding another protein of the present invention is set forth in SEQ ID NO:7 and SEQ ID NO:9, with the coding regions extending from nucleotides 67 to 348 and nucleotides 75 to 356, respectively. These
35 polynucleotides have been identified as "clone H174-10" and "clone H174-43", respectively (collectively referred to herein as "H174"). The amino acid sequence of

5 the protein encoded by clones H174 is set forth in SEQ ID NO:8 and SEQ ID NO:10. Clone H174 was deposited with the American Type Culture Collection on August 11, 1995 and given the accession number ATCC 69882. Clones H174 were isolated from a human activated peripheral blood mononuclear cell (PBMC) library using a trap which selects for nucleotides encoding secreted proteins; therefore, H174 does encode
10 a secreted factor. H174 encodes a novel protein; BLASTN/BLASTX or FASTA searches revealed no exact sequence matches. However, a BLASTX search revealed homology between the H174 protein, human IP-10 (accession M33266) and murine CRG-2 (accession M86820) (species homologs). Based upon these homologies, H174 and these homologous proteins are expected to share at least some activities.

15 The sequence of a polynucleotide encoding another protein of the present invention is set forth in SEQ ID NO:11, with the coding region extending from nucleotides 86 to 544. This polynucleotide has been identified as "B18" The amino acid sequence of the protein encoded by clone B18 is set forth in SEQ ID NO:12. Clone B18 was deposited with the American Type Culture Collection on July 6, 1995
20 and assigned accession number ATCC 69868. Clone B18 was isolated from a human activated peripheral blood mononuclear cell (PBMC) library using a trap which selects for nucleotides encoding secreted proteins; therefore, clone B18 does encode a secreted factor. B18 encodes a novel protein; BLASTN/BLASTX or FASTA searches revealed no exact sequence matches. However, a BLASTX search revealed that the
25 region from amino acid 29 to amino acid 163 of B18 (SEQ ID NO:12) shows marked homology to portions of murine CTLA-8 (amino acids 18 to 150, accession L13839) and herpesvirus *Saimiri* ORF13 ("herpes CTLA-8") (amino acids 19 to 151, accession X64346). Based upon these homologies, B18 is believed to be the human homolog of murine and herpes CTLA-8 (i.e., "human CTLA-8"). B18 may demonstrate proinflammatory activity, particularly in development of T-cell dependent immune
30 responses. B18 is also expected to possess other activities specified herein.

Clones J5, L105, H174 and B18 were each transfected into COS cells labelled with ³⁵S-methionine and protein was expressed. Autoradiographs evidencing expression of the proteins in conditioned media are presented in Figs. 1, 2, 3 and 4,

5 respectively. The bands of protein expressed from the relevant clone are indicated by arrows.

 Polynucleotides hybridizing to the polynucleotides of the present invention under stringent conditions and highly stringent conditions are also part of the present invention. As used herein, "highly stringent conditions" include, for example, at least
10 about 0.2xSSC at 65°C; and "stringent conditions" include, for example, at least about 4xSSC at 65°C or at least about 50% formamide, 4xSSC at 42°C. Allelic variants of the polynucleotides of the present invention are also encompassed by the invention.

 Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention.
15 Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing
20 the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein, such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a decavalent form of the protein
25 of the invention.

 The isolated polynucleotide of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the
30 art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed
35 (transfected) with the ligated polynucleotide/expression control sequence.

5 A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue,
10 primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins.
15 Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments
20 may be accomplished using known chemical or enzymatic methods.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in
25 kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

30 The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (i.e., from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an
35 affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or

5 Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion
10 protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, MA), Pharmacia (Piscataway, NJ) and InVitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed
15 to such epitope. One such epitope ("Flag") is commercially available from Kodak (New Haven, CT).

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the
20 protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic
25 animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are
30 known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic
35 compounds and in immunological processes for the development of antibodies.

5 The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the
10 replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Mutagenic techniques for such replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584).

15 Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are believed to be encompassed by the present invention.

20

USES AND BIOLOGICAL ACTIVITY

 The polynucleotides of the present invention and the proteins encoded thereby are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for
25 proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

RESEARCH TOOL UTILITY

30 The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in
35 disease states); as molecular weight markers on Southern gels; as chromosome markers (when labeled) to map related gene positions; to compare with endogenous

5 DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise
10 anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which
15 binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used to raise antibodies or to elicit another immune response; as a reagent (including the labelled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding
20 protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors
25 of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these "research tool" utilities are capable of being developed into reagent grade or kit format for commercialization as "research products."

30

CYTOKINE AND CELL PROLIFERATION/DIFFERENTIATION ACTIVITY

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting)
35 activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited

5 activity in one or more factor dependent cell proliferation assays, and hence the assays
serve as a convenient confirmation of cytokine activity. The activity of a protein of
the present invention is evidenced by any one of a number of routine factor dependent
cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G,
T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2,
10 CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be
measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those
described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek,
15 D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and
Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-
3.19; Chapter 7, *Immunologic studies in Humans*); Takai et al., *J. Immunol.*
137:3494-3500, 1986; Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990;
Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J.*
20 *Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node
cells or thymocytes include, without limitation, those described in: *Polyclonal T cell*
stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in*
Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons,
25 Toronto. 1994; and *Measurement of mouse and human Interferon γ* , Schreiber, R.D.
In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John
Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and
lymphopoietic cells include, without limitation, those described in: *Measurement of*
30 *Human and Murine Interleukin 2 and Interleukin 4*, Bottomly, K., Davis, L.S. and
Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp.
6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., *J. Exp. Med.*
173:1205-1211, 1991; Moreau et al., *Nature* 336:690-692, 1988; Greenberger et al.,
Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; *Measurement of mouse and human*
35 *interleukin 6* - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds.
Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., *Proc. Natl.*

- 5 Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 -
Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in*
Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto.
1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J.,
Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan
10 eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among
others, proteins that affect APC-T cell interactions as well as direct T-cell effects by
measuring proliferation and cytokine production) include, without limitation, those
described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek,
15 D.H. Margulies, E.M. Shevach, W Strober
Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays
for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors;
Chapter 7, Immunologic studies in Humans); Weinberger et al., *Proc. Natl. Acad. Sci.*
USA 77:6091-6095, 1980; Weinberger et al., *Eur. J. Immun.* 11:405-411, 1981; Takai
20 et al., *J. Immunol.* 137:3494-3500, 1986; Takai et al., *J. Immunol.* 140:508-512,
1988.

IMMUNE STIMULATING/SUPPRESSING ACTIVITY

A protein of the present invention may also exhibit immune stimulating or
25 immune suppressing activity, including without limitation the activities for which
assays are described herein. A protein may be useful in the treatment of various
immune deficiencies and disorders (including severe combined immunodeficiency
(SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B
lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell
30 populations. These immune deficiencies may be genetic or be caused by viral (e.g.,
HIV) as well as bacterial or fungal infections, or may result from autoimmune
disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or
other infection may be treatable using a protein of the present invention, including
infections by HIV, hepatitis viruses, herpes viruses, mycobacteria, leishmania, malaria
35 and various fungal infections such as candida. Of course, in this regard, a protein of

5 the present invention may also be useful where a boost to the immune system generally would be indicated, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome,
10 autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, asthma and related respiratory
15 conditions), may also be treatable using a protein of the present invention.

A protein of the present invention may also suppress chronic or acute inflammation, such as, for example, that associated with infection (such as septic shock or systemic inflammatory response syndrome (SIRS)), inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF
20 or IL-1 (such as the effect demonstrated by IL-11).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan,
25 A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al.,
30 J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular
35 Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

5 Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current*
10 *Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

 Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan,
15 A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

20 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by denritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., *Journal of Experimenal Medicine* 173:549-559, 1991; Macatonia et al., *Journal of Immunology* 154:5071-5079, 1995; Porgador et al., *Journal of Experimental Medicine*
25 182:255-260, 1995; Nair et al., *Journal of Virology* 67:4062-4069, 1993; Huang et al., *Science* 264:961-965, 1994; Macatonia et al., *Journal of Experimental Medicine* 169:1255-1264, 1989; Bhardwaj et al., *Journal of Clinical Investigation* 94:797-807, 1994; and Inaba et al., *Journal of Experimental Medicine* 172:631-640, 1990.

 Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in:
30 Darzynkiewicz et al., *Cytometry* 13:795-808, 1992; Gorczyca et al., *Leukemia* 7:659-670, 1993; Gorczyca et al., *Cancer Research* 53:1945-1951, 1993; Itoh et al., *Cell* 66:233-243, 1991; Zacharchuk, *Journal of Immunology* 145:4037-4045, 1990;
35 Zamai et al., *Cytometry* 14:891-897, 1993; Gorczyca et al., *International Journal of Oncology* 1:639-648, 1992.

5 Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

10

HEMATOPOIESIS REGULATING ACTIVITY

 A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent
15 cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation
20 of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use
25 in place of or complementarily to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post
30 irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e. in conjunction with bone marrow transplantation) as normal cells or genetically manipulated for gene therapy.

 The activity of a protein of the invention may, among other means, be measured by the following methods:

35 Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

- 5 Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.
- 10 Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911,
- 15 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp.
- 20 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162,
- 25 Wiley-Liss, Inc., New York, NY. 1994.

TISSUE GENERATION/REGENERATION ACTIVITY

A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair, and in the treatment of burns, incisions and ulcers.

A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints.

5 *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

10 A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by
15 inflammatory processes.

20 Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the present
25 invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include
30 an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.
35

5 The protein of the present invention may also be useful for proliferation of
neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of
central and peripheral nervous system diseases and neuropathies, as well as
mechanical and traumatic disorders, which involve degeneration, death or trauma to
neural cells or nerve tissue. More specifically, a protein may be used in the treatment
10 of diseases of the peripheral nervous system, such as peripheral nerve injuries,
peripheral neuropathy and localized neuropathies, and central nervous system
diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic
lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated
in accordance with the present invention include mechanical and traumatic disorders,
15 such as spinal cord disorders, head trauma and cerebrovascular diseases such as
stroke. Peripheral neuropathies resulting from chemotherapy or other medical
therapies may also be treatable using a protein of the invention.

 It is expected that a protein of the present invention may also exhibit activity
for generation of other tissues, such as organs (including, for example, pancreas, liver,
20 intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and
vascular (including vascular endothelium) tissue, or for promoting the growth of cells
comprising such tissues. Part of the desired effects may be by inhibition of fibrotic
scarring to allow normal tissue to regenerate.

 A protein of the present invention may also be useful for gut protection or
regeneration and treatment of lung or liver fibrosis, reperfusion injury in various
25 tissues, and conditions resulting from systemic cytokine damage.

 The activity of a protein of the invention may, among other means, be
measured by the following methods:

 Assays for tissue generation activity include, without limitation, those
30 described in: International Patent Publication No. WO95/16035 (bone, cartilage,
tendon); International Patent Publication No. WO95/05846 (nerve, neuronal);
International Patent Publication No. WO91/07491 (skin, endothelium).

ACTIVIN/INHIBIN ACTIVITY

35 A protein of the present invention may also exhibit activin- or inhibin-related
activities. Inhibins are characterized by their ability to inhibit the release of follicle

5 stimulating hormone (FSH), while activins and are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin α family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration
10 of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- β group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885.
15 A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

20 Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

25 CHEMOTACTIC/CHEMOKINETIC ACTIVITY

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, neutrophils, T-cells, mast cells, eosinophils and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilized or attract a desired
30 cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

35 A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such

5 cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

10 The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current
15 Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867,
20 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

HEMOSTATIC AND THROMBOLYTIC ACTIVITY

A protein of the invention may also exhibit hemostatic or thrombolytic
25 activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of
30 conditions resulting therefrom (such as, for example, infarction or stroke).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al.,
35 Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

5 **RECEPTOR/LIGAND ACTIVITY**

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, 10 receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the 15 relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

20 Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 25 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltzenberg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

OTHER ACTIVITIES

30 A protein of the invention may also exhibit one or more of the following additional activities or effects: killing infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin or other tissue pigmentation, or organ size (such as, for example, breast augmentation 35 or diminution); effecting the processing of dietary fat, protein or carbohydrate; effecting behavioral characteristics, including, without limitation, appetite, libido,

5 stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; and in the case of enzymes, correcting deficiencies of the enzyme and treating related diseases.

10

ADMINISTRATION AND DOSING

A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) 15 diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of 20 administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. The pharmaceutical composition may further contain other 25 agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in formulations of the particular cytokine, lymphokine, other 30 hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, 35 pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

5 The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell
10 receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively
15 antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

 The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other
20 pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within
25 the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

 As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that
30 is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active
35 ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

5 In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines,
10 lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on
15 the appropriate sequence of administering protein of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in
20 a variety of conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet,
25 capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier
30 such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical
35 composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

5 When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art.

10 A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may

15 also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

 The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has

20 undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at

25 that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01 μ g to about 100 mg (preferably about 0.1 μ g to about 10 mg, more preferably about 0.1 μ g to about 1 mg) of protein of the present invention per kg body weight.

30 The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous

35 administration. Ultimately the attending physician will decide on the appropriate

5 duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. Such antibodies may be obtained using either the entire protein or fragments thereof as an
10 immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, J. Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, FEBS Lett. 211, 10 (1987). Monoclonal antibodies binding to the
15 protein of the invention may be useful diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal
20 antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When
25 administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention
30 which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a
35 structure for the developing bone and cartilage and optimally capable of being

5 resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential
10 matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are
15 nonbiodegradable and chemically defined, such as sintered hydroxapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size,
20 particle shape, and biodegradability.

Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein
25 compositions from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose, the most preferred being cationic salts
30 of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the
35 polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby

5 providing the protein the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth
10 factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins of the present invention.

15 The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the
20 severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect the dosage. Progress can be monitored by periodic assessment of
25 tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be
30 administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

35

5 Patent and literature references cited herein are incorporated by reference as
if fully set forth.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Jacobs, Kenneth
McCoy, John
Kelleher, Kerry
Carlin, McKeough
- (ii) TITLE OF INVENTION: DNA SEQUENCES AND SECRETED PROTEINS
ENCODED THEREBY
- (iii) NUMBER OF SEQUENCES: 12
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Genetics Institute, Inc. -- Legal Affairs
 - (B) STREET: 87 CambridgePark Drive
 - (C) CITY: Cambridge
 - (D) STATE: Massachusetts
 - (E) COUNTRY: USA
 - (F) ZIP: 02140
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Brown, Scott A.
 - (B) REGISTRATION NUMBER: 32,724
 - (C) REFERENCE/DOCKET NUMBER: GI6000
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (617) 498-8224
 - (B) TELEFAX: (617) 876-5851

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 2209 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 38..1447

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAGAAGATAA AACTGGACAC TGGGGAGACA CAACTTC ATG CTG CGT GGG ATC TCC	55
Met Leu Arg Gly Ile Ser	
1 5	
CAG CTA CCT GCA GTG GCC ACC ATG TCT TGG GTC CTG CTG CCT GTA CTT	103
Gln Leu Pro Ala Val Ala Thr Met Ser Trp Val Leu Leu Pro Val Leu	
10 15 20	
TGG CTC ATT GTT CAA ACT CAA GCA ATA GCC ATA AAG CAA ACA CCT GAA	151
Trp Leu Ile Val Gln Thr Gln Ala Ile Ala Ile Lys Gln Thr Pro Glu	
25 30 35	
TTA ACG CTC CAT GAA ATA GTT TGT CCT AAA AAA CTT CAC ATT TTA CAC	199
Leu Thr Leu His Glu Ile Val Cys Pro Lys Lys Leu His Ile Leu His	
40 45 50	
AAA AGA GAG ATC AAG AAC AAC CAG ACA GAA AAG CAT GGC AAA GAG GAA	247
Lys Arg Glu Ile Lys Asn Asn Gln Thr Glu Lys His Gly Lys Glu Glu	
55 60 65 70	
AGG TAT GAA CCT GAA GTT CAA TAT CAG ATG ATC TTA AAT GGA GAA GAA	295
Arg Tyr Glu Pro Glu Val Gln Tyr Gln Met Ile Leu Asn Gly Glu Glu	
75 80 85	
ATC ATT CTC TCC CTA CAA AAA ACC AAG CAC CTC CTG GGG CCA GAC TAC	343
Ile Ile Leu Ser Leu Gln Lys Thr Lys His Leu Leu Gly Pro Asp Tyr	
90 95 100	
ACT GAA ACA TTG TAC TCA CCC AGA GGA GAG GAA ATT ACC ACG AAA CCT	391
Thr Glu Thr Leu Tyr Ser Pro Arg Gly Glu Glu Ile Thr Thr Lys Pro	
105 110 115	
GAG AAC ATG GAA CAC TGT TAC TAT AAA GGA AAC ATC CTA AAT GAA AAG	439
Glu Asn Met Glu His Cys Tyr Tyr Lys Gly Asn Ile Leu Asn Glu Lys	
120 125 130	
AAT TCT GTT GCC AGC ATC AGT ACT TGT GAC GGG TTG AGA GGA TAC TTC	487
Asn Ser Val Ala Ser Ile Ser Thr Cys Asp Gly Leu Arg Gly Tyr Phe	
135 140 145 150	
ACA CAT CAT CAC CAA AGA TAC CAG ATA AAA CCT CTG AAA AGC ACA GAC	535
Thr His His His Gln Arg Tyr Gln Ile Lys Pro Leu Lys Ser Thr Asp	
155 160 165	
GAG AAA GAA CAT GCC GTC TTT ACA TCT AAC CAG GAG GAA CAA GAC CCA	583
Glu Lys Glu His Ala Val Phe Thr Ser Asn Gln Glu Glu Gln Asp Pro	
170 175 180	
GCT AAC CAC ACA TGT GGT GTG AAG AGC ACT GAC GGG AAA CAA GGC CCA	631
Ala Asn His Thr Cys Gly Val Lys Ser Thr Asp Gly Lys Gln Gly Pro	
185 190 195	
ATT CGA ATC TCT AGA TCA CTC AAA AGC CCA GAG AAA GAA GAC TTT CTT	679
Ile Arg Ile Ser Arg Ser Leu Lys Ser Pro Glu Lys Glu Asp Phe Leu	
200 205 210	
CGG GCA CAG AAA TAC ATT GAT CTC TAT TTG GTG CTG GAT AAT GCC TTT	727
Arg Ala Gln Lys Tyr Ile Asp Leu Tyr Leu Val Leu Asp Asn Ala Phe	
215 220 225 230	
TAT AAG AAC TAT AAT GAG AAT CTA ACT CTG ATA AGA AGC TTT GTG TTT	775

Tyr Lys Asn Tyr Asn Glu Asn Leu Thr Leu Ile Arg Ser Phe Val Phe
235 240 245

GAT GTG ATG AAC CTA CTC AAT GTG ATA TAT AAC ACC ATA GAT GTT CAA Asp Val Met Asn Leu Leu Asn Val Ile Tyr Asn Thr Ile Asp Val Gln 250 255 260	823
GTG GCC TTG GTA GGT ATG GAA ATC TGG TCT GAT GGG GAT AAG ATA AAG Val Ala Leu Val Gly Met Glu Ile Trp Ser Asp Gly Asp Lys Ile Lys 265 270 275	871
GTG GTG CCC AGC GCA AGC ACC ACG TTT GAC AAC TTC CTG AGA TGG CAC Val Val Pro Ser Ala Ser Thr Thr Phe Asp Asn Phe Leu Arg Trp His 280 285 290	919
AGT TCT AAC CTG GGG AAA AAG ATC CAC GAC CAT GCT CAG CTT CTC AGC Ser Ser Asn Leu Gly Lys Lys Ile His Asp His Ala Gln Leu Leu Ser 295 300 305 310	967
GGG ATT AGC TTC AAC AAT CGA CGT GTG GGA CTG GCA GCT TCA AAT TCC Gly Ile Ser Phe Asn Asn Arg Arg Val Gly Leu Ala Ala Ser Asn Ser 315 320 325	1015
TTG TGT TCC CCA TCT TCG GTT GCT GTT ATT GAG GCT AAA AAA AAG AAT Leu Cys Ser Pro Ser Ser Val Ala Val Ile Glu Ala Lys Lys Lys Asn 330 335 340	1063
AAT GTG GCT CTT GTA GGA GTG ATG TCA CAT GAG CTG GGC CAT GTC CTT Asn Val Ala Leu Val Gly Val Met Ser His Glu Leu Gly His Val Leu 345 350 355	1111
GGT ATG CCT GAT GTT CCA TTC AAC ACC AAG TGT CCC TCT GGC AGT TGT Gly Met Pro Asp Val Pro Phe Asn Thr Lys Cys Pro Ser Gly Ser Cys 360 365 370	1159
GTG ATG AAT CAG TAT CTG AGT TCA AAA TTC CCA AAG GAT TTC AGT ACA Val Met Asn Gln Tyr Leu Ser Ser Lys Phe Pro Lys Asp Phe Ser Thr 375 380 385 390	1207
TCT TGC CGT GCA CAT TTT GAA AGA TAC CTT TTA TCT CAG AAA CCA AAG Ser Cys Arg Ala His Phe Glu Arg Tyr Leu Leu Ser Gln Lys Pro Lys 395 400 405	1255
TGC CTG CTG CAA GCA CCT ATT CCT ACA AAT ATA ATG ACA ACA CCA GTG Cys Leu Leu Gln Ala Pro Ile Pro Thr Asn Ile Met Thr Thr Pro Val 410 415 420	1303
TGT GGG AAC CAC CTT CTA GAA GTG GGA GAA GAC TGT GAT TGT GGC TCT Cys Gly Asn His Leu Leu Glu Val Gly Glu Asp Cys Asp Cys Gly Ser 425 430 435	1351
CCT AAG GAG TGT ACC AAT CTC TGC TGT GAA GCC CTA ACG TGT AAA CTG Pro Lys Glu Cys Thr Asn Leu Cys Cys Glu Ala Leu Thr Cys Lys Leu 440 445 450	1399
AAG CCT GGA ACT GAT TGC GGA GGA GAT GCT CCA AAC CAT ACC ACA GAG Lys Pro Gly Thr Asp Cys Gly Gly Asp Ala Pro Asn His Thr Thr Glu 455 460 465 470	1447
TGAATCCAAA AGTCTGCTTC ACTGAGATGC TACCTTGCCA GGACAAGAAC CAAGAACTCT	1507
AACTGTCCCA GGAATCTTGT GAATTTTCAC CCATAATGGT CTTTCACTTG TCATTCTACT	1567
TTCTATATTG TTATCAGTCC AGGAAACAGG TAAACAGATG TAATTAGAGA CATTGGCTCT	1627
TTGTTTAGGC CTAATCTTTC TTTTACTTTT TTTTTTCTT TTTTCTTTT TTTTAAAGAT	1687

```

CATGAATTTG TGACTTAGTT CTGCCCTTTG GAGAACAAAA GAAAGCAGTC TTCCATCAAA 1747
TCACCTTAAA ATGCACGGCT AAACATTTCA GAGTTAACAC TCCAGAATTG TTAAATTACA 1807
AGTACTATGC TTTAATGCTT CTTTCATCTT ACTAGTATGG CCTATAAAAA AAATAATACC 1867
ACTTGATGGG TGAAGGCTTT GGCAATAGAA AGAAGAATAG AATTCAGGTT TTATGTTATT 1927
CCTCTGTGTT CACTTCGCCT TGCTCTTGAA AGTGCAGTAT TTTTCTACAT CATGTCGAGA 1987
ATGATTCAAT GTAAATATTT TTCATTTTAT CATGTATATC CTATACACAC ATCTCCTTCA 2047
TCATCATATA TGAAGTTTAT TTTGAGAAGT CTACATTGCT TACATTTTAA TTGAGCCAGC 2107
AAAGAAGGCT TAATGATTTA TTGAACCATA ATGTCAATAA AAACACAAC TTTGAGGCAA 2167
AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AA 2209

```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 470 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Leu Arg Gly Ile Ser Gln Leu Pro Ala Val Ala Thr Met Ser Trp
 1           5           10           15
Val Leu Leu Pro Val Leu Trp Leu Ile Val Gln Thr Gln Ala Ile Ala
          20           25           30
Ile Lys Gln Thr Pro Glu Leu Thr Leu His Glu Ile Val Cys Pro Lys
          35           40           45
Lys Leu His Ile Leu His Lys Arg Glu Ile Lys Asn Asn Gln Thr Glu
          50           55           60
Lys His Gly Lys Glu Glu Arg Tyr Glu Pro Glu Val Gln Tyr Gln Met
          65           70           75           80
Ile Leu Asn Gly Glu Glu Ile Ile Leu Ser Leu Gln Lys Thr Lys His
          85           90           95
Leu Leu Gly Pro Asp Tyr Thr Glu Thr Leu Tyr Ser Pro Arg Gly Glu
          100          105          110
Glu Ile Thr Thr Lys Pro Glu Asn Met Glu His Cys Tyr Tyr Lys Gly
          115          120          125
Asn Ile Leu Asn Glu Lys Asn Ser Val Ala Ser Ile Ser Thr Cys Asp
          130          135          140
Gly Leu Arg Gly Tyr Phe Thr His His His Gln Arg Tyr Gln Ile Lys
          145          150          155          160
Pro Leu Lys Ser Thr Asp Glu Lys Glu His Ala Val Phe Thr Ser Asn
          165          170          175

```

Gln Glu Glu Gln Asp Pro Ala Asn His Thr Cys Gly Val Lys Ser Thr
 180 185 190
 Asp Gly Lys Gln Gly Pro Ile Arg Ile Ser Arg Ser Leu Lys Ser Pro
 195 200 205
 Glu Lys Glu Asp Phe Leu Arg Ala Gln Lys Tyr Ile Asp Leu Tyr Leu
 210 215 220
 Val Leu Asp Asn Ala Phe Tyr Lys Asn Tyr Asn Glu Asn Leu Thr Leu
 225 230 235 240
 Ile Arg Ser Phe Val Phe Asp Val Met Asn Leu Leu Asn Val Ile Tyr
 245 250 255
 Asn Thr Ile Asp Val Gln Val Ala Leu Val Gly Met Glu Ile Trp Ser
 260 265 270
 Asp Gly Asp Lys Ile Lys Val Val Pro Ser Ala Ser Thr Thr Phe Asp
 275 280 285
 Asn Phe Leu Arg Trp His Ser Ser Asn Leu Gly Lys Lys Ile His Asp
 290 295 300
 His Ala Gln Leu Leu Ser Gly Ile Ser Phe Asn Asn Arg Arg Val Gly
 305 310 315 320
 Leu Ala Ala Ser Asn Ser Leu Cys Ser Pro Ser Ser Val Ala Val Ile
 325 330 335
 Glu Ala Lys Lys Lys Asn Asn Val Ala Leu Val Gly Val Met Ser His
 340 345 350
 Glu Leu Gly His Val Leu Gly Met Pro Asp Val Pro Phe Asn Thr Lys
 355 360 365
 Cys Pro Ser Gly Ser Cys Val Met Asn Gln Tyr Leu Ser Ser Lys Phe
 370 375 380
 Pro Lys Asp Phe Ser Thr Ser Cys Arg Ala His Phe Glu Arg Tyr Leu
 385 390 395 400
 Leu Ser Gln Lys Pro Lys Cys Leu Leu Gln Ala Pro Ile Pro Thr Asn
 405 410 415
 Ile Met Thr Thr Pro Val Cys Gly Asn His Leu Leu Glu Val Gly Glu
 420 425 430
 Asp Cys Asp Cys Gly Ser Pro Lys Glu Cys Thr Asn Leu Cys Cys Glu
 435 440 445
 Ala Leu Thr Cys Lys Leu Lys Pro Gly Thr Asp Cys Gly Gly Asp Ala
 450 455 460
 Pro Asn His Thr Thr Glu
 465 470

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2582 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 52..2034

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATTTCTCAGC TCCAAGCATT AGGTAAACCC ACCAAGCAAT CCTAGCCTGT G ATG GCG	57
Met Ala	
1	
TTT GAC GTC AGC TGC TTC TTT TGG GTG GTG CTG TTT TCT GCC GGC TGT	105
Phe Asp Val Ser Cys Phe Phe Trp Val Val Leu Phe Ser Ala Gly Cys	
5 10 15	
AAA GTC ATC ACC TCC TGG GAT CAG ATG TGC ATT GAG AAA GAA GCC AAC	153
Lys Val Ile Thr Ser Trp Asp Gln Met Cys Ile Glu Lys Glu Ala Asn	
20 25 30	
AAA ACA TAT AAC TGT GAA AAT TTA GGT CTC AGT GAA ATC CCT GAC ACT	201
Lys Thr Tyr Asn Cys Glu Asn Leu Gly Leu Ser Glu Ile Pro Asp Thr	
35 40 45 50	
CTA CCA AAC ACA ACA GAA TTT TTG GAA TTC AGC TTT AAT TTT TTG CCT	249
Leu Pro Asn Thr Thr Glu Phe Leu Glu Phe Ser Phe Asn Phe Leu Pro	
55 60 65	
ACA ATT CAC AAT AGA ACC TTC AGC AGA CTC ATG AAT CTT ACC TTT TTG	297
Thr Ile His Asn Arg Thr Phe Ser Arg Leu Met Asn Leu Thr Phe Leu	
70 75 80	
GAT TTA ACT AGG TGC CAG ATT AAC TGG ATA CAT GAA GAC ACT TTT CAA	345
Asp Leu Thr Arg Cys Gln Ile Asn Trp Ile His Glu Asp Thr Phe Gln	
85 90 95	
AGC CAT CAT CAA TTA AGC ACA CTT GTG TTA ACT GGA AAT CCC CTG ATA	393
Ser His His Gln Leu Ser Thr Leu Val Leu Thr Gly Asn Pro Leu Ile	
100 105 110	
TTC ATG GCA GAA ACA TCG CTT AAT GGG CCC AAG TCA CTG AAG CAT CTT	441
Phe Met Ala Glu Thr Ser Leu Asn Gly Pro Lys Ser Leu Lys His Leu	
115 120 125 130	
TTC TTA ATC CAA ACG GGA ATA TCC AAT CTC GAG TTT ATT CCA GTG CAC	489
Phe Leu Ile Gln Thr Gly Ile Ser Asn Leu Glu Phe Ile Pro Val His	
135 140 145	
AAT CTG GAA AAC TTG GAA AGC TTG TAT CTT GGA AGC AAC CAT ATT TCC	537
Asn Leu Glu Asn Leu Glu Ser Leu Tyr Leu Gly Ser Asn His Ile Ser	
150 155 160	
TCC ATT AAG TTC CCC AAA GAC TTC CCA GCA CGG AAT CTG AAA GTA CTG	585
Ser Ile Lys Phe Pro Lys Asp Phe Pro Ala Arg Asn Leu Lys Val Leu	
165 170 175	
GAT TTT CAG AAT AAT GCT ATA CAC TAC ATC TCT AGA GAA GAC ATG AGG	633

Asp Phe Gln Asn Asn Ala Ile His Tyr Ile Ser Arg Glu Asp Met Arg	
180 185 190	
TCT CTG GAG CAG GCC ATC AAC CTA AGC CTG AAC TTC AAT GGC AAT AAT	681
Ser Leu Glu Gln Ala Ile Asn Leu Ser Leu Asn Phe Asn Gly Asn Asn	
195 200 205 210	
GTT AAA GGT ATT GAG CTT GGG GCT TTT GAT TCA ACG GTC TTC CAA AGT	729
Val Lys Gly Ile Glu Leu Gly Ala Phe Asp Ser Thr Val Phe Gln Ser	
215 220 225	
TTG AAC TTT GGA GGA ACT CCA AAT TTG TCT GTT ATA TTC AAT GGT CTG	777
Leu Asn Phe Gly Thr Pro Asn Leu Ser Val Ile Phe Asn Gly Leu	
230 235 240	
CAG AAC TCT ACT ACT CAG TCT CTC TGG CTG GGA ACA TTT GAG GAC ATT	825
Gln Asn Ser Thr Thr Gln Ser Leu Trp Leu Gly Thr Phe Glu Asp Ile	
245 250 255	
GAT GAC GAA GAT ATT AGT TCA GCC ATG CTC AAG GGA CTC TGT GAA ATG	873
Asp Asp Glu Asp Ile Ser Ser Ala Met Leu Lys Gly Leu Cys Glu Met	
260 265 270	
TCT GTT GAG AGC CTC AAC CTG CAG GAA CAC CGC TTC TCT GAC ATC TCA	921
Ser Val Glu Ser Leu Asn Leu Gln Glu His Arg Phe Ser Asp Ile Ser	
275 280 285 290	
TCC ACC ACA TTT CAG TGC TTC ACC CAA CTC CAA GAA TTG GAT CTG ACA	969
Ser Thr Thr Phe Gln Cys Phe Thr Gln Leu Gln Glu Leu Asp Leu Thr	
295 300 305	
GCA ACT CAC TTG AAA GGG TTA CCC TCT GGG ATG AAG GGT CTG AAC TTG	1017
Ala Thr His Leu Lys Gly Leu Pro Ser Gly Met Lys Gly Leu Asn Leu	
310 315 320	
CTC AAG AAA TTA GTT CTC AGT GTA AAT CAT TTC GAT CAA TTG TGT CAA	1065
Leu Lys Lys Leu Val Leu Ser Val Asn His Phe Asp Gln Leu Cys Gln	
325 330 335	
ATC AGT GCT GCC AAT TTC CCC TCC CTT ACA CAC CTC TAC ATC AGA GGC	1113
Ile Ser Ala Ala Asn Phe Pro Ser Leu Thr His Leu Tyr Ile Arg Gly	
340 345 350	
AAC GTG AAG AAA CTT CAC CTT GGT GTT GGC TGC TTG GAG AAA CTA GGA	1161
Asn Val Lys Lys Leu His Leu Gly Val Gly Cys Leu Glu Lys Leu Gly	
355 360 365 370	
AAC CTT CAG ACA CTT GAT TTA AGC CAT AAT GAC ATA GAG GCT TCT GAC	1209
Asn Leu Gln Thr Leu Asp Leu Ser His Asn Asp Ile Glu Ala Ser Asp	
375 380 385	
TGC TGC AGT CTG CAA CTC AAA AAC CTG TCC CAC TTG CAA ACC TTA AAC	1257
Cys Cys Ser Leu Gln Leu Lys Asn Leu Ser His Leu Gln Thr Leu Asn	
390 395 400	
CTG AGC CAC AAT GAG CCT CTT GGT CTC CAG AGT CAG GCA TTC AAA GAA	1305
Leu Ser His Asn Glu Pro Leu Gly Leu Gln Ser Gln Ala Phe Lys Glu	
405 410 415	
TGT CCT CAG CTA GAA CTC CTC GAT TTG GCA TTT ACC CGC TTA CAC ATT	1353
Cys Pro Gln Leu Glu Leu Leu Asp Leu Ala Phe Thr Arg Leu His Ile	
420 425 430	

AAT GCT CCA CAA AGT CCC TTC CAA AAC CTC CAT TTC CTT CAG GTT CTG Asn Ala Pro Gln Ser Pro Phe Gln Asn Leu His Phe Leu Gln Val Leu 435 440 445 450	1401
AAT CTC ACT TAC TGC TTC CTT GAT ACC AGC AAT CAG CAT CTT CTA GCA Asn Leu Thr Tyr Cys Phe Leu Asp Thr Ser Asn Gln His Leu Leu Ala 455 460 465	1449
GGC CTA CCA GTT CTC CGG CAT CTC AAC TTA AAA GGG AAT CAC TTT CAA Gly Leu Pro Val Leu Arg His Leu Asn Leu Lys Gly Asn His Phe Gln 470 475 480	1497
GAT GGG ACT ATC ACG AAG ACC AAC CTA CTT CAG ACC GTG GGC AGC TTG Asp Gly Thr Ile Thr Lys Thr Asn Leu Leu Gln Thr Val Gly Ser Leu 485 490 495	1545
GAG GTT CTG ATT TTG TCC TCT TGT GGT CTC CTC TCT ATA GAC CAG CAA Glu Val Leu Ile Leu Ser Ser Cys Gly Leu Leu Ser Ile Asp Gln Gln 500 505 510	1593
GCA TTC CAC AGC TTG GGA AAA ATG AGC CAT GTA GAC TTA AGC CAC AAC Ala Phe His Ser Leu Gly Lys Met Ser His Val Asp Leu Ser His Asn 515 520 525 530	1641
AGC CTG ACA TGC GAC AGC ATT GAT TCT CTT AGC CAT CTT AAG GGA ATC Ser Leu Thr Cys Asp Ser Ile Asp Ser Leu Ser His Leu Lys Gly Ile 535 540 545	1689
TAC CTC AAT CTG GCT GCC AAC AGC ATT AAC ATC ATC TCA CCC CGT CTC Tyr Leu Asn Leu Ala Ala Asn Ser Ile Asn Ile Ile Ser Pro Arg Leu 550 555 560	1737
CTC CCT ATC TTG TCC CAG CAG AGC ACC ATT AAT TTA AGT CAT AAC CCC Leu Pro Ile Leu Ser Gln Gln Ser Thr Ile Asn Leu Ser His Asn Pro 565 570 575	1785
CTG GAC TGC ACT TGC TCG AAT ATT CAT TTC TTA ACA TGG TAC AAA GAA Leu Asp Cys Thr Cys Ser Asn Ile His Phe Leu Thr Trp Tyr Lys Glu 580 585 590	1833
AAC CTG CAC AAA CTT GAA GGC TCG GAG GAG ACC ACG TGT GCA AAC CCG Asn Leu His Lys Leu Glu Gly Ser Glu Glu Thr Thr Cys Ala Asn Pro 595 600 605 610	1881
CCA TCT CTA AGG GGA GTT AAG CTA TCT GAT GTC AAG CTT TCC TGT GGG Pro Ser Leu Arg Gly Val Lys Leu Ser Asp Val Lys Leu Ser Cys Gly 615 620 625	1929
ATT ACA GCC ATA GGC ATT TTC TTT CTC ATA GTA TTT CTA TTA TTG TTG Ile Thr Ala Ile Gly Ile Phe Phe Leu Ile Val Phe Leu Leu Leu Leu 630 635 640	1977
GCT ATT CTG CTA TTT TTT GCA GTT AAA TAC CTT CTC AGG TGG AAA TAC Ala Ile Leu Phe Phe Ala Val Lys Tyr Leu Leu Arg Trp Lys Tyr 645 650 655	2025
CAA CAC ATT TAGTGCTGAA GGTTTCCAGA GAAAGCAAAT AAGTGTGCTT Gln His Ile 660	2074
AGCAAAATTG CTCTAAGTGA AAGAACTGTC ATCTGCTGGT GACCAGACCA GACTTTTCAG	2134
ATTGCTTCCT GGAAGTGGC AGGGACTCAC TGTGCTTTTC TGAGCTTCTT ACTCCTGTGA	2194

```

GTCCCAGAGC TAAAGAACCT TCTAGGCAAG TACACCGAAT GACTCAGTCC AGAGGGTCAG      2254
ATGCTGCTGT GAGAGGCACA GAGCCCTTTC CGCATGTGGA AGAGTGGGAG GAAGCAGAGG      2314
GAGGGACTGG GCAGGGACTG CCGGCCCCCG AGTCTCCAC AGGGAGGCCA TTCCCTTCT      2374
ACTCACCGAC ATCCCTCCCA GCACCACACA CCCC GCCCCT GAAAGGAGAT CATCAGCCCC      2434
CACAATTTGT CAGAGCTGAA GCCAGCCCAC TACCCACCCC CACTACAGCA TTGTGCTTGG      2494
GTCTGGGTTT TCAGTAATGT AGCCATTGTA GAAACTTACT TGGGGACAAA GTCTCAATCC      2554
TTATTTTAAA TGAAAAA AAAA AAAA      2582

```

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 661 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Ala Phe Asp Val Ser Cys Phe Phe Trp Val Val Leu Phe Ser Ala
 1             5             10             15
Gly Cys Lys Val Ile Thr Ser Trp Asp Gln Met Cys Ile Glu Lys Glu
 20             25             30
Ala Asn Lys Thr Tyr Asn Cys Glu Asn Leu Gly Leu Ser Glu Ile Pro
 35             40             45
Asp Thr Leu Pro Asn Thr Thr Glu Phe Leu Glu Phe Ser Phe Asn Phe
 50             55             60
Leu Pro Thr Ile His Asn Arg Thr Phe Ser Arg Leu Met Asn Leu Thr
 65             70             75             80
Phe Leu Asp Leu Thr Arg Cys Gln Ile Asn Trp Ile His Glu Asp Thr
 85             90             95
Phe Gln Ser His His Gln Leu Ser Thr Leu Val Leu Thr Gly Asn Pro
 100            105            110
Leu Ile Phe Met Ala Glu Thr Ser Leu Asn Gly Pro Lys Ser Leu Lys
 115            120            125
His Leu Phe Leu Ile Gln Thr Gly Ile Ser Asn Leu Glu Phe Ile Pro
 130            135            140
Val His Asn Leu Glu Asn Leu Glu Ser Leu Tyr Leu Gly Ser Asn His
 145            150            155            160
Ile Ser Ser Ile Lys Phe Pro Lys Asp Phe Pro Ala Arg Asn Leu Lys
 165            170            175
Val Leu Asp Phe Gln Asn Asn Ala Ile His Tyr Ile Ser Arg Glu Asp
 180            185            190
Met Arg Ser Leu Glu Gln Ala Ile Asn Leu Ser Leu Asn Phe Asn Gly

```

195					200					205					
Asn	Asn	Val	Lys	Gly	Ile	Glu	Leu	Gly	Ala	Phe	Asp	Ser	Thr	Val	Phe
210						215					220				
Gln	Ser	Leu	Asn	Phe	Gly	Gly	Thr	Pro	Asn	Leu	Ser	Val	Ile	Phe	Asn
225					230					235					240
Gly	Leu	Gln	Asn	Ser	Thr	Thr	Gln	Ser	Leu	Trp	Leu	Gly	Thr	Phe	Glu
				245					250					255	
Asp	Ile	Asp	Asp	Glu	Asp	Ile	Ser	Ser	Ala	Met	Leu	Lys	Gly	Leu	Cys
			260					265					270		
Glu	Met	Ser	Val	Glu	Ser	Leu	Asn	Leu	Gln	Glu	His	Arg	Phe	Ser	Asp
		275					280					285			
Ile	Ser	Ser	Thr	Thr	Phe	Gln	Cys	Phe	Thr	Gln	Leu	Gln	Glu	Leu	Asp
	290					295					300				
Leu	Thr	Ala	Thr	His	Leu	Lys	Gly	Leu	Pro	Ser	Gly	Met	Lys	Gly	Leu
305					310					315					320
Asn	Leu	Leu	Lys	Lys	Leu	Val	Leu	Ser	Val	Asn	His	Phe	Asp	Gln	Leu
				325					330					335	
Cys	Gln	Ile	Ser	Ala	Ala	Asn	Phe	Pro	Ser	Leu	Thr	His	Leu	Tyr	Ile
			340					345					350		
Arg	Gly	Asn	Val	Lys	Lys	Leu	His	Leu	Gly	Val	Gly	Cys	Leu	Glu	Lys
		355					360					365			
Leu	Gly	Asn	Leu	Gln	Thr	Leu	Asp	Leu	Ser	His	Asn	Asp	Ile	Glu	Ala
	370					375					380				
Ser	Asp	Cys	Cys	Ser	Leu	Gln	Leu	Lys	Asn	Leu	Ser	His	Leu	Gln	Thr
385					390					395					400
Leu	Asn	Leu	Ser	His	Asn	Glu	Pro	Leu	Gly	Leu	Gln	Ser	Gln	Ala	Phe
				405					410					415	
Lys	Glu	Cys	Pro	Gln	Leu	Glu	Leu	Leu	Asp	Leu	Ala	Phe	Thr	Arg	Leu
			420					425					430		
His	Ile	Asn	Ala	Pro	Gln	Ser	Pro	Phe	Gln	Asn	Leu	His	Phe	Leu	Gln
		435					440					445			
Val	Leu	Asn	Leu	Thr	Tyr	Cys	Phe	Leu	Asp	Thr	Ser	Asn	Gln	His	Leu
	450					455					460				
Leu	Ala	Gly	Leu	Pro	Val	Leu	Arg	His	Leu	Asn	Leu	Lys	Gly	Asn	His
465					470					475					480
Phe	Gln	Asp	Gly	Thr	Ile	Thr	Lys	Thr	Asn	Leu	Leu	Gln	Thr	Val	Gly
				485					490					495	
Ser	Leu	Glu	Val	Leu	Ile	Leu	Ser	Ser	Cys	Gly	Leu	Leu	Ser	Ile	Asp
			500					505					510		
Gln	Gln	Ala	Phe	His	Ser	Leu	Gly	Lys	Met	Ser	His	Val	Asp	Leu	Ser
		515					520					525			
His	Asn	Ser	Leu	Thr	Cys	Asp	Ser	Ile	Asp	Ser	Leu	Ser	His	Leu	Lys

530 535 540
 Gly Ile Tyr Leu Asn Leu Ala Ala Asn Ser Ile Asn Ile Ile Ser Pro
 545 550 555 560
 Arg Leu Leu Pro Ile Leu Ser Gln Gln Ser Thr Ile Asn Leu Ser His
 565 570 575
 Asn Pro Leu Asp Cys Thr Cys Ser Asn Ile His Phe Leu Thr Trp Tyr
 580 585 590
 Lys Glu Asn Leu His Lys Leu Glu Gly Ser Glu Glu Thr Thr Cys Ala
 595 600 605
 Asn Pro Pro Ser Leu Arg Gly Val Lys Leu Ser Asp Val Lys Leu Ser
 610 615 620
 Cys Gly Ile Thr Ala Ile Gly Ile Phe Phe Leu Ile Val Phe Leu Leu
 625 630 635 640
 Leu Leu Ala Ile Leu Leu Phe Phe Ala Val Lys Tyr Leu Leu Arg Trp
 645 650 655
 Lys Tyr Gln His Ile
 660

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 588 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 76..474

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CGGCCAAAGA GGCCTAAACT TGC GGCTGTC CATCTCACCT ACAGCTCTGG TCTCATCCTC 60
 AACTCAACCA CAATC ATG GCT CAG ATG ATG ACT CTG AGC CTC CTT AGC CTG 111
 Met Ala Gln Met Met Thr Leu Ser Leu Leu Ser Leu
 1 5 10
 GTC CTG GCT CTC TGC ATC CCC TGG ACC CAA GGC AGT GAT GGA GGG GGT 159
 Val Leu Ala Leu Cys Ile Pro Trp Thr Gln Gly Ser Asp Gly Gly Gly
 15 20 25
 CAG GAC TGC TGC CTT AAG TAC AGC CAG AAG AAA ATT CCC TAC AGT ATT 207
 Gln Asp Cys Cys Leu Lys Tyr Ser Gln Lys Lys Ile Pro Tyr Ser Ile
 30 35 40
 GTC CGA GGC TAT AGG AAG CAA GAA CCA AGT TTA GGC TGT CCC ATC CCG 255
 Val Arg Gly Tyr Arg Lys Gln Glu Pro Ser Leu Gly Cys Pro Ile Pro
 45 50 55 60

GCA ATC CTG TTC TCA CCC CGG AAG CAC TCT AAG CCT GAG CTA TGT GCA 303
 Ala Ile Leu Phe Ser Pro Arg Lys His Ser Lys Pro Glu Leu Cys Ala
 65 70 75

AAC CCT GAG GAA GGC TGG GTG CAG AAC CTG ATG CGC CGC CTG GAC CAG 351
 Asn Pro Glu Glu Gly Trp Val Gln Asn Leu Met Arg Arg Leu Asp Gln
 80 85 90

CCT CCA GCC CCA GGG AAA CAA AGC CCC GGC TGC AGG AAG AAC CGG GGA 399
 Pro Pro Ala Pro Gly Lys Gln Ser Pro Gly Cys Arg Lys Asn Arg Gly
 95 100 105

ACC TCT AAG TCT GGA AAG AAA GGA AAG GGC TCC AAG GGC TGC AAG AGA 447
 Thr Ser Lys Ser Gly Lys Lys Gly Lys Gly Ser Lys Gly Cys Lys Arg
 110 115 120

ACT GAA CAG ACA CAG CCC TCA AGA GGA TAGCCAGTA GCCCGCCTGG 494
 Thr Glu Gln Thr Gln Pro Ser Arg Gly
 125 130

AGCCCAGGAG ATCCCCCAGC AACTTCAAGC TGGGTGGTTC ACGGTCCAAC TCACAGGCAA 554

AGAGGGAGCT AGAAAACAGA CTCAGGAGCC GCTA 588

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 133 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Gln Met Met Thr Leu Ser Leu Leu Ser Leu Val Leu Ala Leu
 1 5 10 15

Cys Ile Pro Trp Thr Gln Gly Ser Asp Gly Gly Gly Gln Asp Cys Cys
 20 25 30

Leu Lys Tyr Ser Gln Lys Lys Ile Pro Tyr Ser Ile Val Arg Gly Tyr
 35 40 45

Arg Lys Gln Glu Pro Ser Leu Gly Cys Pro Ile Pro Ala Ile Leu Phe
 50 55 60

Ser Pro Arg Lys His Ser Lys Pro Glu Leu Cys Ala Asn Pro Glu Glu
 65 70 75 80

Gly Trp Val Gln Asn Leu Met Arg Arg Leu Asp Gln Pro Pro Ala Pro
 85 90 95

Gly Lys Gln Ser Pro Gly Cys Arg Lys Asn Arg Gly Thr Ser Lys Ser
 100 105 110

Gly Lys Lys Gly Lys Gly Ser Lys Gly Cys Lys Arg Thr Glu Gln Thr
 115 120 125

Gln Pro Ser Arg Gly
 130

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 966 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 67..348

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

CTTCCAAGAA GAGCAGCAAA GCTGAAGTAG CAGCAACAGC ACCAGCAGCA ACAGCAAAAA      60
ACAAAC ATG AGT GTG AAG GGC ATG GCT ATA GCC TTG GCT GTG ATA TTG      108
    Met Ser Val Lys Gly Met Ala Ile Ala Leu Ala Val Ile Leu
        1             5             10

TGT GCT ACA GTT GTT CAA GGC TTC CCC ATG TTC AAA AGA GGA CGC TGT      156
Cys Ala Thr Val Val Gln Gly Phe Pro Met Phe Lys Arg Gly Arg Cys
    15             20             25             30

CTT TGC ATA GGC CCT GGG GTA AAA GCA GTG AAA GTG GCA GAT ATT GAG      204
Leu Cys Ile Gly Pro Gly Val Lys Ala Val Lys Val Ala Asp Ile Glu
        35             40             45

AAA GCC TCC ATA ATG TAC CCA AGT AAC AAC TGT GAC AAA ATA GAA GTG      252
Lys Ala Ser Ile Met Tyr Pro Ser Asn Asn Cys Asp Lys Ile Glu Val
        50             55             60

ATT ATT ACC CTG AAA GAA AAT AAA GGA CAA CGA TGC CTA AAT CCC AAA      300
Ile Ile Thr Leu Lys Glu Asn Lys Gly Gln Arg Cys Leu Asn Pro Lys
        65             70             75

TCG AAG CAA GCA AGG CTT ATA ATC AAA AAA GTT GAA AGA AAG AAT TTT      348
Ser Lys Gln Ala Arg Leu Ile Ile Lys Lys Val Glu Arg Lys Asn Phe
    80             85             90

TAAAAATATC AAAACATATG AAGTCTTGGA AAAGGGCATC TGAAAAACCT AGAACAAGTT      408
TAACTGTGAC TACTGAAATG ACAAGAATTC TACAGTAGGA AACTGAGACT TTTCTATGGT      468
TTTGTGACTT TCAACTTTTG TACAGTTATG TGAAGGATGA AAGGTGGGTG AAAGGACCAA      528
AAACAGAAAT ACAGTCTTCC TGAATGAATG ACAATCAGAA TTCCACTGCC CAAAGGAGTC      588
CAACAATTAA ATGGATTTCT AGGAAAAGCT ACCTTAAGAA AGGCTGGTTA CCATCGGAGT      648
TTACAAAGTG CTTTCACGTT CTTACTTGTT GTATTATACA TTCATGCATT TCTAGGCTAG      708
AGAACCTTCT AGATTTGATG CTTACAATA TTCTGTTGTG ACTATGAGAA CATTTCTGTC      768
TCTAGAAGTT ATCTGTCTGT ATTGATCTTT ATGCTATATT ACTATCTGTG GTTACAGTGG      828
AGACATTGAC ATTATTACTG GAGTCAAGCC CTTATAAGTC AAAAGCACCT ATGTGTCGTA      888

```


AAGCATTTCCT CAAACATTTA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 948
 AAAAAAAAAA AAAAAAAAAA 966

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 94 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Ser Val Lys Gly Met Ala Ile Ala Leu Ala Val Ile Leu Cys Ala 15
 1 5 10
 Thr Val Val Gln Gly Phe Pro Met Phe Lys Arg Gly Arg Cys Leu Cys 30
 20 25 30
 Ile Gly Pro Gly Val Lys Ala Val Lys Val Ala Asp Ile Glu Lys Ala 45
 35 40 45
 Ser Ile Met Tyr Pro Ser Asn Asn Cys Asp Lys Ile Glu Val Ile Ile 60
 50 55 60
 Thr Leu Lys Glu Asn Lys Gly Gln Arg Cys Leu Asn Pro Lys Ser Lys 80
 65 70 75 80
 Gln Ala Arg Leu Ile Ile Lys Lys Val Glu Arg Lys Asn Phe 90
 85 90

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1354 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 75..356

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TTCTACTCCT TCCAAGAAGA GCAGCAAAGC TGAAGTAGCA GCAACAGCAC CAGCAGCAAC 60
 AGCAAAAAAC AAAC ATG AGT GTG AAG GGC ATG GCT ATA GCC TTG GCT GTG 110
 Met Ser Val Lys Gly Met Ala Ile Ala Leu Ala Val 10
 1 5 10
 ATA TTG TGT GCT ACA GTT GTT CAA GGC TTC CCC ATG TTC AAA AGA GGA 158
 Ile Leu Cys Ala Thr Val Val Gln Gly Phe Pro Met Phe Lys Arg Gly 25
 15 20 25

CGC TGT CTT TGC ATA GGC CCT GGG GTA AAA GCA GTG AAA GTG GCA GAT	206
Arg Cys Leu Cys Ile Gly Pro Gly Val Lys Ala Val Lys Val Ala Asp	
30 35 40	
ATT GAG AAA GCC TCC ATA ATG TAC CCA AGT AAC AAC TGT GAC AAA ATA	254
Ile Glu Lys Ala Ser Ile Met Tyr Pro Ser Asn Asn Cys Asp Lys Ile	
45 50 55 60	
GAA GTG ATT ATT ACC CTG AAA GAA AAT AAA GGA CAA CGA TGC CTA AAT	302
Glu Val Ile Ile Thr Leu Lys Glu Asn Lys Gly Gln Arg Cys Leu Asn	
65 70 75	
CCC AAA TCG AAG CAA GCA AGG CTT ATA ATC AAA AAA GTT GAA AGA AAG	350
Pro Lys Ser Lys Gln Ala Arg Leu Ile Ile Lys Lys Val Glu Arg Lys	
80 85 90	
AAT TTT TAAAAATATC AAAACATATG AAGTCCTGGA AAAGGGCATC TGAAAAACCT	406
Asn Phe	
AGAACAAGTT TAACTGTGAC TACTGAAATG ACAAGAATTC TACAGTAGGA AACTGAGACT	466
TTTCTATGGT TTTGTGACTT TCAACTTTTG TACAGTTATG TGAAGGATGA AAGGTGGGTG	526
AAAGGACCAA AAACAGAAAT ACAGTCTTCC TGAATGAATG ACAATCAGAA TTCCACTGCC	586
CAAAGGAGTC CAACAATTAA ATGGATTTCT AGGAAAAGCT ACCTTAAGAA AGGCTGGTTA	646
CCATCGGAGT TTACAAAGTG CTTTCACGTT CTTACTTGTT GTATTATACA TTCATGCATT	706
TCTAGGCTAG AGAACCTTCT AGATTTGATG CTTACAAC TA TCTGTTGTG ACTATGAGAA	766
CATTTCTGTC TCTAGAAGTT ATCTGTCTGT ATTGATCTTT ATGCTATATT ACTATCTGTG	826
GTTACAGTGG AGACATTGAC ATTATTACTG GAGTCAAGCC CTTATAAGTC AAAAGCACCT	886
ATGTGTCGTA AAGCATTCTT CAAACATTTT TTCATGCAAA TACACACTTC TTTCCCCAAA	946
TATCATGTAG CACATCAATA TGTAGGGAAA CATTCTTATG CATCATTTGG TTTGTTTTAT	1006
AACCAATTCA TTAAATGTAA TTCATAAAAT GTACTATGAA AAAAATTATA CGCTATGGGA	1066
TACTGGCAAC AGTGCACATA TTTCATAACC AAATTAGCAG CACCGGTCTT AATTTGATGT	1126
TTTTCAACTT TTATTCATTG AGATGTTTTG AAGCAATTAG GATATGTGTG TTTACTGTAC	1186
TTTTTGTTTT GATCCGTTTG TATAAATGAT AGCAATATCT TGGACACATT TGAAATACAA	1246
AATGTTTTTG TCTACCAAAG AAAAATGTTG AAAAATAAGC AAATGTATAC CTAGCAATCA	1306
CTTTTACTTT TTGTAATTCT GTCTCTTAGA AAAATACATA ATCTAATT	1354

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 94 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Ser Val Lys Gly Met Ala Ile Ala Leu Ala Val Ile Leu Cys Ala
 1 5 10 15
 Thr Val Val Gln Gly Phe Pro Met Phe Lys Arg Gly Arg Cys Leu Cys
 20 25 30
 Ile Gly Pro Gly Val Lys Ala Val Lys Val Ala Asp Ile Glu Lys Ala
 35 40 45
 Ser Ile Met Tyr Pro Ser Asn Asn Cys Asp Lys Ile Glu Val Ile Ile
 50 55 60
 Thr Leu Lys Glu Asn Lys Gly Gln Arg Cys Leu Asn Pro Lys Ser Lys
 65 70 75 80
 Gln Ala Arg Leu Ile Ile Lys Lys Val Glu Arg Lys Asn Phe
 85 90

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 813 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 86..544

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GGGAAGATAC ATTACAGAA AGAGCTTCCT GCACAAAGTA AGCCACCAGC GCAACATGAC 60
 AGTGAAGACC CTGCATGGCC CAGCC ATG GTC AAG TAC TTG CTG CTG TCG ATA 112
 Met Val Lys Tyr Leu Leu Leu Ser Ile
 1 5
 TTG GGG CTT GCC TTT CTG AGT GAG GCG GCA GCT CGG AAA ATC CCC AAA 160
 Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Arg Lys Ile Pro Lys
 10 15 20 25
 GTA GGA CAT ACT TTT TTC CAA AAG CCT GAG AGT TGC CCG CCT GTG CCA 208
 Val Gly His Thr Phe Gln Lys Pro Glu Ser Cys Pro Pro Val Pro
 30 35 40
 GGA GGT AGT ATG AAG CTT GAC ATT GGC ATC ATC AAT GAA AAC CAG CGC 256
 Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu Asn Gln Arg
 45 50 55
 GTT TCC ATG TCA CGT AAC ATC GAG AGC CGC TCC ACC TCC CCC TGG AAT 304
 Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser Pro Trp Asn
 60 65 70
 TAC ACT GTC ACT TGG GAC CCC AAC CGG TAC CCC TCG GAA GTT GTA CAG 352
 Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu Val Val Gln
 75 80 85

GCC CAG TGT AGG AAC TTG GGC TGC ATC AAT GCT CAA GGA AAG GAA GAC 400
 Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly Lys Glu Asp
 90 95 100 105
 ATC TCC ATG AAT TCC GTT CCC ATC CAG CAA GAG ACC CTG GTC GTC CGG 448
 Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu Val Val Arg
 110 115 120
 AGG AAG CAC CAA GGC TGC TCT GTT TCT TTC CAG TTG GAG AAG GTG CTG 496
 Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu Lys Val Leu
 125 130 135
 GTG ACT GTT GGC TGC ACC TGC GTC ACC CCT GTC ATC CAC CAT GTG CAG 544
 Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His His Val Gln
 140 145 150
 TAAGAGGTGC ATATCCACTC AGCTGAAGAA GCTGTAGAAA TGCCACTCCT TACCCAGTGC 604
 TCTGCAACAA GTCCTGTCTG ACCCCCAATT CCCTCCACTT CACAGGACTC TTAATAAGAC 664
 CTGCACGGAT GGAAACAGAA AATATTACACA ATGTATGTGT GTATGTACTA CACTTTATAT 724
 TTGATATCTA AAATGTTAGG AGAAAAATTA ATATATTCAG TGCTAATATA ATAAAGTATT 784
 AATAATTTAA AAATAAAAAA AAAAAAAA 813

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 153 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Val Lys Tyr Leu Leu Leu Ser Ile Leu Gly Leu Ala Phe Leu Ser
 1 5 10 15
 Glu Ala Ala Ala Arg Lys Ile Pro Lys Val Gly His Thr Phe Phe Gln
 20 25 30
 Lys Pro Glu Ser Cys Pro Pro Val Pro Gly Gly Ser Met Lys Leu Asp
 35 40 45
 Ile Gly Ile Ile Asn Glu Asn Gln Arg Val Ser Met Ser Arg Asn Ile
 50 55 60
 Glu Ser Arg Ser Thr Ser Pro Trp Asn Tyr Thr Val Thr Trp Asp Pro
 65 70 75 80
 Asn Arg Tyr Pro Ser Glu Val Val Gln Ala Gln Cys Arg Asn Leu Gly
 85 90 95
 Cys Ile Asn Ala Gln Gly Lys Glu Asp Ile Ser Met Asn Ser Val Pro
 100 105 110
 Ile Gln Gln Glu Thr Leu Val Val Arg Arg Lys His Gln Gly Cys Ser
 115 120 125
 Val Ser Phe Gln Leu Glu Lys Val Leu Val Thr Val Gly Cys Thr Cys

130

135

140

Val Thr Pro Val Ile His His Val Gln
145 150

What is claimed is:

1. A composition comprising an isolated polynucleotide selected from the group consisting of:
 - (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 38 to nucleotide 1447;
 - (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:1 encoding a protein having biological activity;
 - (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
 - (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity;
 - (e) a polynucleotide which is an allelic variant of SEQ ID NO:1;and
 - (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).
2. A composition of claim 1 wherein said polynucleotide is operably linked to an expression control sequence.
3. A host cell transformed with a composition of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein, which comprises:
 - (a) growing a culture of the host cell of claim 3 in a suitable culture medium; and
 - (b) purifying the protein from the culture
6. A protein produced according to the process of claim 5.
7. A composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2; and
 - (b) fragments of the amino acid sequence of SEQ ID NO:2;
- the protein being substantially free from other mammalian proteins.

8. The composition of claim 7, further comprising a pharmaceutically acceptable carrier.

9. A composition comprising an antibody which specifically reacts with the protein of claim 7.

10. A method for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition of claim 8.

11. A composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 52 to nucleotide 2034;
- (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:3 encoding a protein having biological activity;
- (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity;
- (e) a polynucleotide which is an allelic variant of SEQ ID NO:4; and
- (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

12. A composition of claim 11 wherein said polynucleotide is operably linked to an expression control sequence.

13. A host cell transformed with a composition of claim 12.
14. The host cell of claim 13, wherein said cell is a mammalian cell.
15. A process for producing a protein, which comprises:
 - (a) growing a culture of the host cell of claim 13 in a suitable culture medium; and
 - (b) purifying the protein from the culture
16. A protein produced according to the process of claim 15.
17. A composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:
 - (a) the amino acid sequence of SEQ ID NO:4; and
 - (b) fragments of the amino acid sequence of SEQ ID NO:4;the protein being substantially free from other mammalian proteins.
18. The composition of claim 17, further comprising a pharmaceutically acceptable carrier.
19. A composition comprising an antibody which specifically reacts with the protein of claim 17.
20. A method for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition of claim 18.
21. A composition comprising an isolated polynucleotide selected from the group consisting of:
 - (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:5 from nucleotide 76 to nucleotide 474;

- (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:5 encoding a protein having biological activity;
- (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:6;
- (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity;
- (e) a polynucleotide which is an allelic variant of SEQ ID NO:5;
- and
- (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

22. A composition of claim 21 wherein said polynucleotide is operably linked to an expression control sequence.

23. A host cell transformed with a composition of claim 22.

24. The host cell of claim 23, wherein said cell is a mammalian cell.

25. A process for producing a protein, which comprises:

- (a) growing a culture of the host cell of claim 23 in a suitable culture medium; and
- (b) purifying the protein from the culture

26. A protein produced according to the process of claim 25.

27. A composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6; and
 - (b) fragments of the amino acid sequence of SEQ ID NO:6;
- the protein being substantially free from other mammalian proteins.

28. The composition of claim 27, further comprising a pharmaceutically acceptable carrier.

29. A composition comprising an antibody which specifically reacts with the protein of claim 27.

30. A method for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition of claim 28.

31. A composition comprising an isolated polynucleotide selected from the group consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 67 to nucleotide 348;

(b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:7 encoding a protein having biological activity;

(c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:8;

(d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity;

(e) a polynucleotide which is an allelic variant of SEQ ID NO:7; and

(f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

32. A composition of claim 31 wherein said polynucleotide is operably linked to an expression control sequence.

33. A host cell transformed with a composition of claim 32.

34. The host cell of claim 33, wherein said cell is a mammalian cell.

35. A process for producing a protein, which comprises:
- (a) growing a culture of the host cell of claim 33 in a suitable culture medium; and
 - (b) purifying the protein from the culture
36. A protein produced according to the process of claim 35.
37. A composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:8; and
 - (b) fragments of the amino acid sequence of SEQ ID NO:8;
- the protein being substantially free from other mammalian proteins.
38. The composition of claim 37, further comprising a pharmaceutically acceptable carrier.
39. A composition comprising an antibody which specifically reacts with the protein of claim 37.
40. A method for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition of claim 38.
41. A composition comprising an isolated polynucleotide selected from the group consisting of:
- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 75 to nucleotide 356;
 - (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:9 encoding a protein having biological activity;
 - (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;

- (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity;
- (e) a polynucleotide which is an allelic variant of SEQ ID NO:9;
- and
- (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

42. A composition of claim 41 wherein said polynucleotide is operably linked to an expression control sequence.

43. A host cell transformed with a composition of claim 42.

44. The host cell of claim 43, wherein said cell is a mammalian cell.

45. A process for producing a protein, which comprises:

- (a) growing a culture of the host cell of claim 43 in a suitable culture medium; and
- (b) purifying the protein from the culture

46. A protein produced according to the process of claim 45.

47. A composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10; and
 - (b) fragments of the amino acid sequence of SEQ ID NO:10;
- the protein being substantially free from other mammalian proteins.

48. The composition of claim 47, further comprising a pharmaceutically acceptable carrier.

49. A composition comprising an antibody which specifically reacts with the protein of claim 47.

50. A method for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition of claim 48.

51. A composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 86 to nucleotide 544;
- (b) a polynucleotide comprising a fragment of the nucleotide sequence of SEQ ID NO:11 encoding a protein having biological activity;
- (c) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;
- (d) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity;
- (e) a polynucleotide which is an allelic variant of SEQ ID NO:11; and
- (f) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(e).

52. A composition of claim 51 wherein said polynucleotide is operably linked to an expression control sequence.

53. A host cell transformed with a composition of claim 52.

54. The host cell of claim 53, wherein said cell is a mammalian cell.

55. A process for producing a protein, which comprises:

- (a) growing a culture of the host cell of claim 53 in a suitable culture medium; and
- (b) purifying the protein from the culture

56. A protein produced according to the process of claim 55.

57. A composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12; and
- (b) fragments of the amino acid sequence of SEQ ID NO:12;

the protein being substantially free from other mammalian proteins.

58. The composition of claim 57, further comprising a pharmaceutically acceptable carrier.

59. A composition comprising an antibody which specifically reacts with the protein of claim 57.

60. A method for preventing, treating or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of a composition of claim 58.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.